

## XXXIV. Sampling from a Normal Population

---

### Setting

- We have a population of stuff, e.g. students of different heights.
  - The population has a mean  $\mu$  and variance  $\sigma^2$ .
  - We will take samples:  $Y_1 :=$  height of one student.  $Y_2 :=$  height of another, and so on.  $Y_n :=$  height of last sample student.
- 

### Assumptions and Notation

- **Assumption forever:** Our samples are independent.
  - We say the  $Y_i$  are independent identically distributed (i.i.d.) random variables.
  - **Assumption for this lecture only:** The population is normally distributed.
  - **Notation:** We say  $Y_i \sim N(\mu, \sigma^2)$ .
- 

### The Sample Mean

- **Statistic we'll study:** The sample mean  
 $\bar{Y} := \frac{1}{n} (Y_1 + \cdots + Y_n)$ .

- **Theorem:** Suppose  $Y_i \sim N(\mu, \sigma^2)$ . Then

$$\bar{Y} \sim N\left(\mu, \frac{\sigma^2}{n}\right).$$

- That is,  $\bar{Y}$  is normally distributed with mean  $\mu$  and variance  $\frac{\sigma^2}{n}$ .

### Standard Normal Distribution

		Second decimal place of $z$									
$z$	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09	
0.0	.5000	.4960	.4920	.4880	.4840	.4801	.4761	.4721	.4681	.4641	
0.1	.4602	.4562	.4522	.4483	.4443	.4404	.4364	.4325	.4286	.4247	
0.2	.4207	.4168	.4129	.4090	.4052	.4013	.3974	.3936	.3897	.3859	
0.3	.3821	.3783	.3745	.3707	.3669	.3632	.3594	.3557	.3520	.3483	
0.4	.3446	.3409	.3372	.3336	.3300	.3264	.3228	.3192	.3156	.3121	
0.5	.3085	.3050	.3015	.2981	.2946	.2912	.2877	.2843	.2810	.2776	
0.6	.2743	.2709	.2676	.2643	.2611	.2578	.2546	.2514	.2483	.2451	
0.7	.2420	.2389	.2358	.2327	.2296	.2266	.2236	.2206	.2177	.2148	
0.8	.2119	.2090	.2061	.2033	.2005	.1977	.1949	.1922	.1894	.1867	
0.9	.1841	.1814	.1788	.1762	.1736	.1711	.1685	.1660	.1635	.1611	
1.0	.1587	.1562	.1539	.1515	.1492	.1469	.1446	.1423	.1401	.1379	
1.1	.1357	.1335	.1314	.1292	.1271	.1251	.1230	.1210	.1190	.1170	
1.2	.1151	.1131	.1112	.1093	.1075	.1056	.1038	.1020	.1003	.0985	
1.3	.0968	.0951	.0934	.0918	.0901	.0885	.0869	.0853	.0838	.0823	
1.4	.0808	.0793	.0778	.0764	.0749	.0735	.0722	.0708	.0694	.0681	
1.5	.0668	.0655	.0643	.0630	.0618	.0606	.0594	.0582	.0571	.0559	
1.6	.0548	.0537	.0526	.0516	.0505	.0495	.0485	.0475	.0465	.0455	
1.7	.0446	.0436	.0427	.0418	.0409	.0401	.0392	.0384	.0375	.0367	
1.8	.0359	.0352	.0344	.0336	.0329	.0322	.0314	.0307	.0301	.0294	
1.9	.0287	.0281	.0274	.0268	.0262	.0256	.0250	.0244	.0239	.0233	
2.0	.0228	.0222	.0217	.0212	.0207	.0202	.0197	.0192	.0188	.0183	

### Converting to Standard Normal

- **Recall:** If  $\bar{Y}$  is normal, then

$$Z := \frac{\bar{Y} - \text{mean}}{\text{standard deviation}}$$

is standard normal, that is,  $Z \sim N(0, 1)$ .

- **Corollary to Theorem:**

$$Z := \frac{\sqrt{n}(\bar{Y} - \mu)}{\sigma} \sim N(0, 1)$$

- So we can use charts to find probabilities for  $Z$  and then work back to find them for  $Y$ .

---

### Example I

Heights of students are normally distributed with a standard deviation of 4 inches. You measure the heights of 9 students. What is the probability that their mean is within 2 inches of the (unknown) global mean?

$n = 9$ ,  $\mu = \text{unknown}$ ,  $\sigma = 4$ .  $Z := \frac{3(\bar{Y} - \mu)}{4}$  is standard normal.

$$|\bar{Y} - \mu| \leq 2 \text{ iff } \left| \frac{3(\bar{Y} - \mu)}{4} \right| \leq 1.5$$

From the table on page 848,  $P(Z > 1.5) \approx 0.0668$ .  
[Graph the normal curve and shade the tail.]

$$P(Z < -1.5 \text{ or } Z > 1.5) \approx 2 * 0.0668 = 0.1336$$

$$P(-1.5 \leq Z \leq 1.5) \approx 1 - 0.1336 = 0.8664 \approx \boxed{87\%}$$

---

z	Second decimal place of z									
	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
0.0	.5000	.4960	.4920	.4880	.4840	.4801	.4761	.4721	.4681	.4641
0.1	.4602	.4562	.4522	.4483	.4443	.4404	.4364	.4325	.4286	.4247
0.2	.4207	.4168	.4129	.4090	.4052	.4013	.3974	.3936	.3897	.3859
0.3	.3821	.3783	.3745	.3707	.3669	.3632	.3594	.3557	.3520	.3483
0.4	.3446	.3409	.3372	.3336	.3300	.3264	.3228	.3192	.3156	.3121
0.5	.3085	.3050	.3015	.2981	.2946	.2912	.2877	.2843	.2810	.2776
0.6	.2743	.2709	.2676	.2643	.2611	.2578	.2546	.2514	.2483	.2451
0.7	.2420	.2389	.2358	.2327	.2296	.2266	.2236	.2206	.2177	.2148
0.8	.2119	.2090	.2061	.2033	.2005	.1977	.1949	.1922	.1894	.1867
0.9	.1841	.1814	.1788	.1762	.1736	.1711	.1685	.1660	.1635	.1611
1.0	.1587	.1562	.1539	.1515	.1492	.1469	.1446	.1423	.1401	.1379
1.1	.1357	.1335	.1314	.1292	.1271	.1251	.1230	.1210	.1190	.1170
1.2	.1151	.1131	.1112	.1093	.1075	.1056	.1038	.1020	.1003	.0985
1.3	.0968	.0951	.0934	.0918	.0901	.0885	.0869	.0853	.0838	.0823
1.4	.0808	.0793	.0778	.0764	.0749	.0735	.0722	.0708	.0694	.0681
1.5	.0668	.0655	.0643	.0630	.0618	.0606	.0594	.0582	.0571	.0559
1.6	.0548	.0537	.0526	.0516	.0505	.0495	.0485	.0475	.0465	.0455
1.7	.0446	.0436	.0427	.0418	.0409	.0401	.0392	.0384	.0375	.0367
1.8	.0359	.0352	.0344	.0336	.0329	.0322	.0314	.0307	.0301	.0294
1.9	.0287	.0281	.0274	.0268	.0262	.0256	.0250	.0244	.0239	.0233
2.0	.0228	.0222	.0217	.0212	.0207	.0202	.0197	.0192	.0188	.0183

### Example II

Let  $Y_1, \dots, Y_n$  be i.i.d. variables, with  $Y_i \sim N(\mu, \sigma^2)$ . If  $\sigma^2 = 64$  and  $n = 36$ , find the probability that  $\bar{Y}$  is within one unit of  $\mu$ . What happens to this probability as  $n \rightarrow \infty$ ?

$$\frac{\bar{Y} - \mu}{\frac{\sigma^2}{n}} = \frac{\bar{Y} - \mu}{\frac{8}{6}} = \frac{3}{4}(\bar{Y} - \mu)$$

$$|\bar{Y} - \mu| \leq 1 \text{ iff } \frac{3}{4}|\bar{Y} - \mu| \leq \frac{3}{4}$$

$$P \approx 1 - 2(0.2266) = 1 - 0.4532 = \boxed{0.5468 \approx 55\%}$$

This  $\boxed{\rightarrow 1}$  as  $n \rightarrow \infty$ .

z	Second decimal place of z									
	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
0.0	.5000	.4960	.4920	.4880	.4840	.4801	.4761	.4721	.4681	.4641
0.1	.4602	.4562	.4522	.4483	.4443	.4404	.4364	.4325	.4286	.4247
0.2	.4207	.4168	.4129	.4090	.4052	.4013	.3974	.3936	.3897	.3859
0.3	.3821	.3783	.3745	.3707	.3669	.3632	.3594	.3557	.3520	.3483
0.4	.3446	.3409	.3372	.3336	.3300	.3264	.3228	.3192	.3156	.3121
0.5	.3085	.3050	.3015	.2981	.2946	.2912	.2877	.2843	.2810	.2776
0.6	.2743	.2709	.2676	.2643	.2611	.2578	.2546	.2514	.2483	.2451
0.7	.2420	.2389	.2358	.2327	.2296	.2266	.2236	.2206	.2177	.2148
0.8	.2119	.2090	.2061	.2033	.2005	.1977	.1949	.1922	.1894	.1867
0.9	.1841	.1814	.1788	.1762	.1736	.1711	.1685	.1660	.1635	.1611
1.0	.1587	.1562	.1539	.1515	.1492	.1469	.1446	.1423	.1401	.1379
1.1	.1357	.1335	.1314	.1292	.1271	.1251	.1230	.1210	.1190	.1170
1.2	.1151	.1131	.1112	.1093	.1075	.1056	.1038	.1020	.1003	.0985
1.3	.0968	.0951	.0934	.0918	.0901	.0885	.0869	.0853	.0838	.0823
1.4	.0808	.0793	.0778	.0764	.0749	.0735	.0722	.0708	.0694	.0681
1.5	.0668	.0655	.0643	.0630	.0618	.0606	.0594	.0582	.0571	.0559
1.6	.0548	.0537	.0526	.0516	.0505	.0495	.0485	.0475	.0465	.0455
1.7	.0446	.0436	.0427	.0418	.0409	.0401	.0392	.0384	.0375	.0367
1.8	.0359	.0352	.0344	.0336	.0329	.0322	.0314	.0307	.0301	.0294
1.9	.0287	.0281	.0274	.0268	.0262	.0256	.0250	.0244	.0239	.0233
2.0	.0228	.0222	.0217	.0212	.0207	.0202	.0197	.0192	.0188	.0183

### Example III

Students at a university have taken an average of 70 units, with a standard deviation of 20 units. Assuming this distribution is normal, if you sample 9 students, what is the chance that their average unit total is between 67 and 73 units?

We want  $P(|\bar{Y} - \mu| \leq 3)$ . We know

$$Z := \frac{\sqrt{n}(\bar{Y} - \mu)}{\sigma} \sim N(0, 1).$$

$$\sqrt{n} = 3, \sigma = 20$$

$$\begin{aligned} |\bar{Y} - \mu| &\leq 3 \\ \frac{3|\bar{Y} - \mu|}{20} &\leq \frac{3 \cdot 3}{20} = 0.45 \\ P(Z > 0.45) &= 0.3264 \end{aligned}$$

$$P(-0.45 < Z < 0.45) = 1 - 2(0.3264) = 1 - 0.6528 = \boxed{0.3472 \approx 35\%}$$

Second decimal place of $z$										
$z$	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
0.0	.5000	.4960	.4920	.4880	.4840	.4801	.4761	.4721	.4681	.4641
0.1	.4602	.4562	.4522	.4483	.4443	.4404	.4364	.4325	.4286	.4247
0.2	.4207	.4168	.4129	.4090	.4052	.4013	.3974	.3936	.3897	.3859
0.3	.3821	.3783	.3745	.3707	.3669	.3632	.3594	.3557	.3520	.3483
0.4	.3446	.3409	.3372	.3336	.3300	.3264	.3228	.3192	.3156	.3121
0.5	.3085	.3050	.3015	.2981	.2946	.2912	.2877	.2843	.2810	.2776
0.6	.2743	.2709	.2676	.2643	.2611	.2578	.2546	.2514	.2483	.2451
0.7	.2420	.2389	.2358	.2327	.2296	.2266	.2236	.2206	.2177	.2148
0.8	.2119	.2090	.2061	.2033	.2005	.1977	.1949	.1922	.1894	.1867
0.9	.1841	.1814	.1788	.1762	.1736	.1711	.1685	.1660	.1635	.1611
1.0	.1587	.1562	.1539	.1515	.1492	.1469	.1446	.1423	.1401	.1379
1.1	.1357	.1335	.1314	.1292	.1271	.1251	.1230	.1210	.1190	.1170
1.2	.1151	.1131	.1112	.1093	.1075	.1056	.1038	.1020	.1003	.0985
1.3	.0968	.0951	.0934	.0918	.0901	.0885	.0869	.0853	.0838	.0823
1.4	.0808	.0793	.0778	.0764	.0749	.0735	.0722	.0708	.0694	.0681
1.5	.0668	.0655	.0643	.0630	.0618	.0606	.0594	.0582	.0571	.0559
1.6	.0548	.0537	.0526	.0516	.0505	.0495	.0485	.0475	.0465	.0455
1.7	.0446	.0436	.0427	.0418	.0409	.0401	.0392	.0384	.0375	.0367
1.8	.0359	.0352	.0344	.0336	.0329	.0322	.0314	.0307	.0301	.0294
1.9	.0287	.0281	.0274	.0268	.0262	.0256	.0250	.0244	.0239	.0233
2.0	.0228	.0222	.0217	.0212	.0207	.0202	.0197	.0192	.0188	.0183

### Example IV

Six samples are taken from a normally distributed population with variance 0.67. What is the probability that the sample mean will be within 0.5 units of the population mean?

We want  $P(|\bar{Y} - \mu| \leq 0.5)$ . We know

$$Z := \frac{\sqrt{n}(\bar{Y} - \mu)}{\sigma} \sim N(0, 1).$$

$$\begin{aligned} |\bar{Y} - \mu| &\leq 0.5 \\ \frac{\sqrt{6}|\bar{Y} - \mu|}{\sqrt{0.67}} &\leq \frac{\sqrt{6} \cdot 0.5}{\sqrt{0.67}} = \sqrt{9}(0.5) = \frac{3}{2} \\ P(Z > 1.5) &= 0.0668 \end{aligned}$$

$$P(-1.5 < Z < 1.5) = 1 - 2(0.0668) = 1 - 0.1336 = \boxed{0.8664 \approx 87\%}$$

---

z	Second decimal place of z									
	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
0.0	.5000	.4960	.4920	.4880	.4840	.4801	.4761	.4721	.4681	.4641
0.1	.4602	.4562	.4522	.4483	.4443	.4404	.4364	.4325	.4286	.4247
0.2	.4207	.4168	.4129	.4090	.4052	.4013	.3974	.3936	.3897	.3859
0.3	.3821	.3783	.3745	.3707	.3669	.3632	.3594	.3557	.3520	.3483
0.4	.3446	.3409	.3372	.3336	.3300	.3264	.3228	.3192	.3156	.3121
0.5	.3085	.3050	.3015	.2981	.2946	.2912	.2877	.2843	.2810	.2776
0.6	.2743	.2709	.2676	.2643	.2611	.2578	.2546	.2514	.2483	.2451
0.7	.2420	.2389	.2358	.2327	.2296	.2266	.2236	.2206	.2177	.2148
0.8	.2119	.2090	.2061	.2033	.2005	.1977	.1949	.1922	.1894	.1867
0.9	.1841	.1814	.1788	.1762	.1736	.1711	.1685	.1660	.1635	.1611
1.0	.1587	.1562	.1539	.1515	.1492	.1469	.1446	.1423	.1401	.1379
1.1	.1357	.1335	.1314	.1292	.1271	.1251	.1230	.1210	.1190	.1170
1.2	.1151	.1131	.1112	.1093	.1075	.1056	.1038	.1020	.1003	.0985
1.3	.0968	.0951	.0934	.0918	.0901	.0885	.0869	.0853	.0838	.0823
1.4	.0808	.0793	.0778	.0764	.0749	.0735	.0722	.0708	.0694	.0681
1.5	.0668	.0655	.0643	.0630	.0618	.0606	.0594	.0582	.0571	.0559
1.6	.0548	.0537	.0526	.0516	.0505	.0495	.0485	.0475	.0465	.0455
1.7	.0446	.0436	.0427	.0418	.0409	.0401	.0392	.0384	.0375	.0367
1.8	.0359	.0352	.0344	.0336	.0329	.0322	.0314	.0307	.0301	.0294
1.9	.0287	.0281	.0274	.0268	.0262	.0256	.0250	.0244	.0239	.0233
2.0	.0228	.0222	.0217	.0212	.0207	.0202	.0197	.0192	.0188	.0183

### Example V

As in Example IV, we plan to sample a normally distributed population with variance 0.67. We want to ensure that our sample mean will be within 0.5 units of the population mean with probability 95%. How many samples should we take?



We want  $P(|\bar{Y} - \mu| \leq 0.5) = 0.95$ . We know

$$Z := \frac{\sqrt{n}(\bar{Y} - \mu)}{\sigma} \sim N(0, 1).$$

Draw the normal curve. We want 95% of the area in the middle, so we want the outer area to be  $\frac{1 - 0.95}{2} = 0.025$ . This corresponds by the chart to  $Z = 1.96$ .

$$\begin{aligned} |\bar{Y} - \mu| &\leq 0.5 \\ \left| \frac{\sqrt{n}(\bar{Y} - \mu)}{\sigma} \right| &\leq \frac{0.5\sqrt{n}}{\sigma} \\ |Z| &\leq \frac{0.5\sqrt{n}}{\sigma} \\ 1.96 &\leq \frac{0.5\sqrt{n}}{\sigma} \\ \frac{1.96\sigma}{0.5} &\leq \sqrt{n} \\ 3.92\sigma &\leq \sqrt{n} \\ (3.92)(\sqrt{0.67}) &\leq \sqrt{n} \\ n &\geq \left[ (3.92)(\sqrt{0.67}) \right]^2 \approx 10.296 \\ n &= \boxed{11} \end{aligned}$$


---

*Will Murray's Probability, XXXIV. Sampling from a Normal Population* 10

z	Second decimal place of z									
	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
0.0	.5000	.4960	.4920	.4880	.4840	.4801	.4761	.4721	.4681	.4641
0.1	.4602	.4562	.4522	.4483	.4443	.4404	.4364	.4325	.4286	.4247
0.2	.4207	.4168	.4129	.4090	.4052	.4013	.3974	.3936	.3897	.3859
0.3	.3821	.3783	.3745	.3707	.3669	.3632	.3594	.3557	.3520	.3483
0.4	.3446	.3409	.3372	.3336	.3300	.3264	.3228	.3192	.3156	.3121
0.5	.3085	.3050	.3015	.2981	.2946	.2912	.2877	.2843	.2810	.2776
0.6	.2743	.2709	.2676	.2643	.2611	.2578	.2546	.2514	.2483	.2451
0.7	.2420	.2389	.2358	.2327	.2296	.2266	.2236	.2206	.2177	.2148
0.8	.2119	.2090	.2061	.2033	.2005	.1977	.1949	.1922	.1894	.1867
0.9	.1841	.1814	.1788	.1762	.1736	.1711	.1685	.1660	.1635	.1611
1.0	.1587	.1562	.1539	.1515	.1492	.1469	.1446	.1423	.1401	.1379
1.1	.1357	.1335	.1314	.1292	.1271	.1251	.1230	.1210	.1190	.1170
1.2	.1151	.1131	.1112	.1093	.1075	.1056	.1038	.1020	.1003	.0985
1.3	.0968	.0951	.0934	.0918	.0901	.0885	.0869	.0853	.0838	.0823
1.4	.0808	.0793	.0778	.0764	.0749	.0735	.0722	.0708	.0694	.0681
1.5	.0668	.0655	.0643	.0630	.0618	.0606	.0594	.0582	.0571	.0559
1.6	.0548	.0537	.0526	.0516	.0505	.0495	.0485	.0475	.0465	.0455
1.7	.0446	.0436	.0427	.0418	.0409	.0401	.0392	.0384	.0375	.0367
1.8	.0359	.0352	.0344	.0336	.0329	.0322	.0314	.0307	.0301	.0294
1.9	.0287	.0281	.0274	.0268	.0262	.0256	.0250	.0244	.0239	.0233
2.0	.0228	.0222	.0217	.0212	.0207	.0202	.0197	.0192	.0188	.0183